

# Production of Microbial Lipases for the Study of Triglyceride Structure

2356

JOHN A. ALFORD and JAMES L. SMITH, Eastern Utilization Research and Development Division, United States Department of Agriculture, Beltsville, Maryland

## Abstract

Lipases from different microorganisms are known to differ in their site of attack on triglycerides, and recent evidence has indicated that they may be useful in studying triglyceride structure. This paper is concerned with the most recent developments in the production, recovery, and stability of lipases from three of these microorganisms. The lipase from *Staphylococcus aureus*, which attacks both the 1-, 2-, and 3-positions of triglycerides, is produced in an aerated, tryptic digest of casein at 30C in 1-2 days. The lipase from *Geotrichum candidum*, which attacks primarily unsaturated fatty acid linkages and shows some stereospecificity, is produced in a static culture grown on a mineral salts-glucose-protein hydrolysate medium incubated at 20C for 4-5 days. An improved method is described for preparing lyophilized preparations of these, and the *Pseudomonas fragi* lipase, which are quite stable when stored in a refrigerator.

## Introduction

PREVIOUS STUDIES from this laboratory (2,4) have shown that lipases produced by microorganisms attack triglycerides in at least three different ways. The lipase from *Pseudomonas fragi* is similar to pancreatic lipase in that it attacks predominantly the 1-, 3-positions of triglycerides. The lipase from *Staphylococcus aureus* attacks both the 1-, 2-, and 3-positions, while the lipase produced by *Geotrichum candidum* attacks primarily unsaturated fatty acid linkages, regardless of the position of the attachment. More recent evidence has shown that the enzyme from *G. candidum* is stereospecific (5) and, therefore, can be used in studying the structure of triglycerides (6). Because of the interest in these enzymes as possible tools for determining triglyceride structure, particularly the one from *G. candidum*, the present study was undertaken to try to increase their yields, to standardize the recovery procedures, and to examine more closely the stability of the lipases under different conditions of storage.

## Experimental Methods

**Preparation of Media.** The media for all compari-

TABLE I  
Effect of Medium and Aeration on Lipase Production by *Staphylococcus aureus*

Medium	Aerated <sup>a</sup>	Not aerated <sup>b</sup>
	ml <sup>c</sup>	ml
Peptone (Difco)	4.0	1.0
Phytone (BBL)	3.8	1.5
NZ Amine B (Shef.)	4.4	3.7
Trypticase (BBL)	6.0	3.0
Tryptone (Difco)	6.4	1.2
Trypticase soy broth (BBL) <sup>d</sup>	6.0	—
Peptone (Case)	5.5	1.0

<sup>a</sup> Aerated on reciprocating shaker (150-200 cycles/min) at 30C for 1-2 days.

<sup>b</sup> Incubated at 30C for 4 days.

<sup>c</sup> Ml 0.02 N NaOH required for titration of acid released from lard emulsion in 30 min at 35C.

<sup>d</sup> Prepared as described by manufacturer.

sons, except for part of the investigation on *G. candidum*, consisted of: Nitrogen source 1.5% (w/v) and 1M phosphate buffer, 5% (v/v, pH 7.0 unless stated otherwise). For the latter part of the study on *G. candidum*, the salts-glucose-protein hydrolysate medium used was: Protein hydrolysate, 2-4%; NH<sub>4</sub>Cl, 0.1%; KH<sub>2</sub>PO<sub>4</sub>, 0.15%; MgSO<sub>4</sub> · 7H<sub>2</sub>O, 0.012%; and 0.001% of FeSO<sub>4</sub> · 7H<sub>2</sub>O, ZnSO<sub>4</sub> · 7H<sub>2</sub>O, and MnSO<sub>4</sub> · H<sub>2</sub>O. After sterilization the pH was 5.8-6.0, and sterile glucose solution was added to give a final concentration of 0.25%. The sources of the media were as follows: BBL—Baltimore Biological Laboratories, Inc., Baltimore, Md.; Difco—Difco Laboratories, Inc., Detroit, Mich.; Case—Case Laboratories, Inc., Chicago, Ill.; Shef—Sheffield Chemical, Norwich, N.Y.; and NBC—Nutritional Biochemicals Corp., Cleveland, Ohio.

**Assays for Lipase Activity.** The substrates were prepared and the assays for lipolytic activity were carried out as previously described (3).

**Stability Studies.** After the cultures were grown under optimum conditions for lipase production, the cells were removed by centrifugation or, in the case of *G. candidum*, by filtration through coarse filter paper. Portions of this supernatant were stored at 20C and 2C, and samples were removed daily for assay. Additional amounts were shell-frozen in 3-4 ml quantities in test tubes in a dry ice-alcohol bath and stored in a dry ice chest. Tubes of the frozen enzyme were removed at weekly intervals, thawed, and assayed for activity. To avoid any effects of repeated freezing and thawing on the enzyme, none of the excess enzyme was refrozen.

**Lyophilization of Enzymes.** Supernatant solutions obtained as described above were placed in dialysis tubing (1.75 in. flat diameter) in approximately 300-ml quantities. Using the concentration method of Kohn (7), the tubes were placed in a tray, covered with polyethylene glycol (Carbowax 20 M, Union Carbide, N.Y.), and stored at 1-3C for 6-18 hr, during which time the solution was reduced to 10% or less of its original volume. The concentrated solution then was transferred to one or more small dialysis tubes (ca. 50 ml/tube) and dialyzed against at least 100 volumes of cold distilled water (1-3C) for 4-5 hr.

TABLE II  
Comparison of Media for the Production of Lipase by *Geotrichum candidum* at 20C in 4 days

	Titratable acidity <sup>a</sup>
	ml
5% Phosphate buffer (pH 6.0) plus 1.5%:	
Edamin S (Shef.)	1.0
Trypticase (BBL)	4.0
Tryptone (Difco)	3.5
Peptone (Case)	3.2
Peptone (Difco)	3.5
Peptone I (NBC)	1.8
Salts glucose basal medium plus:	
Edamin S (2%)	6.8
Edamin S (4%)	7.3
Trypticase (2%)	6.8
Trypticase (4%)	7.1
Tryptone (2%)	4.8
Peptone (Case) (2%)	5.8

<sup>a</sup> Ml 0.02 N NaOH required for titration of acid released from lard emulsion in 30 min at 35C.

TABLE III

Effect of Temperature and Aeration on Production of Lipase by *Geotrichum candidum*

	Temperature and time of incubation <sup>a</sup>									
	20C			25C				31C		
	Days			Days				Days		ml
	3	4	5	2	3	4	5	2	3	4
Aerated—fast <sup>c</sup>	ml <sup>b</sup>	ml	ml	ml	ml	ml	ml	ml	ml	ml
Aerated—slow <sup>d</sup>	1.9	2.4	3.4	2.1	2.0	2.0	2.0	4.2	6.6	5.5
Not aerated	0.6	7.1	8.2	0.3	8.6	12.0	10.8	3.6	12.5	7.4

<sup>a</sup> 100 ml medium in 1000-ml Erlenmeyer flask.<sup>b</sup> Ml 0.02 N NaOH required for titration of acid released from lard emulsion in 30 min at 35C.<sup>c</sup> Incubated on a rotary shaker at 200 rpm.<sup>d</sup> Incubated on a rotary shaker at 73 rpm.

After dialysis, the contents of the tubes were transferred to suitable flasks, shell-frozen in a dry ice-alcohol bath, and dried under vacuum ( $<20 \mu$  Hg) in a Virtis freeze-dryer. The dry powder was placed in glass vials, stoppered under atmospheric conditions, and stored in a refrigerator at 2–3C.

### Results

**Production of Lipase by *S. aureus*.** In our first work on *S. aureus* (2), a buffered 1% peptone broth was employed; in later work (4), buffered 1% Trypticase (BBL) was used because slightly higher yields were obtained. Statically grown cultures were employed in these earlier studies because aeration had been shown to cause a definite decrease in lipase production by *P. fragi* (1). In the present investigation, the effect of aeration on the production of lipase by *S. aureus* was examined. The data in Table I show that higher yields of lipase could be obtained in a shorter time when the culture was aerated and that the highest yields were produced in the tryptic digest of casein.

**Production of Lipase by *G. candidum*.** The data in Table II show that the highest yields of lipase produced by *G. candidum* were obtained with the salts-glucose medium containing either Edamin Type S (Shef.) or Trypticase (BBL) as the nitrogen source. (Edamin Type S (Shef.) is a lactalbumin hydrolysate.) It should be emphasized that no attempt was made to maintain the pH at 6.0 during growth, and the final pH of the medium was 7.5–8.0. When the initial pH was adjusted to pH 7.0, yields were slightly lower, although the final pH was similar.

The effect of aeration and temperature on the production of *G. candidum* lipase is shown in Table III. The maximum yield of lipase per milliliter of culture was obtained at 31C in 3 days, or at 25C in 4 days in static culture. Destruction of the lipase was faster at the higher temperature, however. Since appreciable lipase was produced with very slow agitation, the effect of increased volumes of culture in flasks of the

TABLE IV

Effect of Surface-to-Volume Ratio of Medium on the Production of Lipase by *Geotrichum candidum*

Volume of medium <sup>a</sup> in 1000-ml Erlenmeyer flask	Method and time of incubation at 25C					
	Nonaerated			Aerated <sup>d</sup>		
	Days			Days		
	3	4	5	3	4	5
ml	ml	ml	ml	ml	ml	ml
100	8.6 <sup>c</sup>	12.0	10.8	7.1	7.0	6.2
250	2.6	2.9	6.2	5.4	7.0	6.7
500	0.4	0.6	0.6	5.1	6.7	6.3

<sup>a</sup> 4% Edamin S-salts-glucose medium.<sup>b</sup> Incubated on a rotary shaker at 73 rpm.<sup>c</sup> Ml NaOH required for titration of acid released from lard emulsion in 30 min at 35C.

same size was determined. The results are shown in Table IV. When the surface-volume ratio of static cultures was decreased, there was a sharp drop in the yield of lipase per milliliter of culture. With the slowly agitated culture, however, yield per milliliter of culture remained essentially unchanged as the volume increased.

**Production of Lipase by *P. fragi*.** The media and cultural conditions for the production of *P. fragi* in both synthetic and partially defined media have been previously described (3). No further increase in yield has been obtained by substituting other protein hydrolysates.

**Stability of Lipase.** The stability of the lipases from the three microorganisms under various conditions of storage is shown in Table V. These data indicate that the lipases are relatively stable, and lyophilized preparations, after an initial loss of up to 50% during processing, can be held for long periods with relatively little change in activity.

### Discussion

By manipulation of the media, temperature, and aeration of the cultures, the production of lipases by *S. aureus* and *G. candidum* has been improved in a manner similar to that previously reported for *P. fragi* (3). It is apparent that the nutritional and oxygen requirements for maximum lipase production vary widely among microorganisms. One must also be aware that, in addition to a varying response depending upon the nitrogen source, variations will occur between products with the same designation (e.g., peptone) but obtained from different manufacturers.

Although the amount of lipase activity per milliliter of medium produced by *G. candidum* in a static culture with a large surface-to-volume ratio was almost twice that obtainable with slow aeration, the total amount of lipase obtainable per flask was nearly three times as great with the aerated culture when the volume of medium per flask was increased. Thus, when large lots of enzyme are to be produced and incuba-

TABLE V

Stability of Lipolytic Enzymes under Different Conditions of Storage

Preparation of enzyme	Loss during preparation			Storage life		
	<i>P. fragi</i>	<i>S. aureus</i>	<i>G. candidum</i>	<i>P. fragi</i>	<i>S. aureus</i>	<i>G. candidum</i>
Supernatant stored at 20C	%	%	%			
Supernatant stored at 2C	N.A. <sup>a</sup>	N.A.	N.A.	2 days <sup>b</sup>	— <sup>c</sup>	3 days <sup>d</sup>
Supernatant frozen; stored in dry ice cabinet	N.A.	N.A.	N.A.	3 days <sup>b</sup>	—	10 days <sup>d</sup>
Lyophilized; stored in refrigerator	20	—	5	>2 mo <sup>e</sup>	—	>2 mo <sup>e</sup>
	25–50	25–50	25–50	>6 mo	>1 yr	>1 yr

<sup>a</sup> Not applicable.<sup>b</sup> Turbidity due to cell growth appeared.<sup>c</sup> Test not done.<sup>d</sup> Time at which approximately 20% of activity was lost.<sup>e</sup> Less than 20% loss of activity by time shown.

tion space is at a premium, use of mild aeration should be considered.

Rapid freezing of the culture supernatant with storage at a low temperature affords a simple procedure for prolonged storage of small quantities of enzyme. However, since the concentration, dialysis, and lyophilization procedure will reduce several liters of material to a few grams, it has obvious advantages for any extensive study.

It is reasonable to expect that further study on the interrelationships of pH, aeration, and supply of nutrients during the entire growth cycle would bring about additional increases in yield of these lipases. However, with the media and conditions of growth de-

scribed here, coupled with proper recovery and storage procedures, stable preparations can be routinely prepared which are suitable for the study of the structure of triglycerides.

## REFERENCES

1. Alford, J. A., and L. E. Elliott, *Food Research* 25, 296-303 (1960).
2. Alford, J. A., and D. A. Pierce, *J. Food Sci.* 26, 518-524 (1961).
3. Alford, J. A., and D. A. Pierce, *J. Bacteriol.* 86, 24-29 (1963).
4. Alford, J. A., D. A. Pierce and F. G. Suggs, *J. Lipid Res.* 5, 390-394 (1964).
5. Jensen, R. G., J. Sampugna, J. G. Quinn, D. L. Carpenter, T. A. Marks and J. A. Alford, *JAOCs* 42, 1029-1032 (1965).
6. Jensen, R. G., J. Sampugna, J. G. Quinn, D. L. Carpenter and J. A. Alford, *J. Dairy Sci.* 48:767 (1965).
7. Kohn, J., *Nature* 183, 1055 (1959).