

[Reprinted from the Journal of Organic Chemistry, 38, 174 (1973).]  
Copyright 1973 by the American Chemical Society and reprinted by permission of the copyright owner.

### Stearoyl Methanesulfonate. A Mixed Anhydride from an Isopropenyl Ester

**Summary:** Isopropenyl stearate and methanesulfonic acid react at ambient temperatures in methylene chloride to give the mixed anhydride stearoyl methanesulfonate, a powerful stearoylating agent.

**Sir:** Mixed carboxylic-oxy acid anhydrides have been regarded as highly effective acylating agents. They have been prepared by the reaction of a silver sulfonate with an acid chloride,<sup>1</sup> a sulfonic acid with a ketene,<sup>2</sup> a sulfonic acid with an acid chloride,<sup>3</sup> or a sulfonic acid with a carboxylic anhydride<sup>3</sup> or from a lithium carboxylate and the sulfur trioxide-*N,N*-dimethylformamide complex.<sup>4</sup>

Nevertheless, little attention has been given to the facile synthesis of these acylating agents from sulfonic acids and isopropenyl esters. An early report<sup>5</sup> on the reactions of isopropenyl acetate (IPA) acknowledged that IPA reacts with carboxylic acids in the presence of a catalytic amount of sulfuric acid to give mixed carboxylic anhydrides. Another group studied the kinetics of various acid-catalyzed IPA reactions and concluded that the active intermediates are mixed acetic-sulfonic or acetic-sulfuric anhydrides.<sup>6</sup> Isopropenyl esters now may be prepared easily by reaction of a carboxylic acid with propyne in the presence of zinc oxide or the zinc carboxylate,<sup>7</sup> and this opens a new synthetic pathway to the once difficult to obtain mixed anhydrides.

We have isolated the mixed anhydride stearoyl methanesulfonate from the reaction mixture of equimolar amounts of isopropenyl stearate and anhydrous methanesulfonic acid in methylene chloride. Although attempts to recrystallize the white solid led to disproportionation, ir analysis and potentiometric titration demonstrated that the initial product was reasonably pure mixed anhydride. Ir absorptions (in a Nujol mull) occur at 1805 and 1185  $\text{cm}^{-1}$  and are typical<sup>10</sup> for mixed carboxylic-sulfonic anhydrides. Ir analysis also showed that the material was essentially free of stearic acid, stearic anhydride, and isopropenyl stearate. A weighed sample of product was hydrolyzed in a known amount of dilute aqueous sodium hydroxide, and the resulting solution was back-titrated with 0.1 *N* hydrochloric acid. Two potentiometric end points were observed, one corresponding to double the amount of sodium hydroxide of the other; one end point was in the pH 5 region (methanesulfonic acid) and the other at pH 9 (stearic acid). The equivalent weight of the mixed anhydride calculated from the titration was within 3% of the theoretical value. The stearic acid that resulted from the hydrolysis and subsequent acidification was isolated in 93% yield and identified by its melting point and its ir spectrum. Quenching of the

mixed anhydride in methanol produced methyl stearate; *tert*-butyl stearate resulted from quenching in *tert*-butyl alcohol.

Gpc evidence confirms that the isolated material that we have labeled "mixed anhydride" is not merely an IPS-methanesulfonic acid adduct  $\text{C}_{17}\text{H}_{35}\text{CO}_2\text{C}(\text{CH}_3)_2\text{OSO}_2\text{CH}_3$ . The gpc trace of a methylene chloride solution of *isolated* mixed anhydride previously quenched with methanol showed the absence of acetone, whereas repetition of the experiment with *in situ* generated mixed anhydride plus methanol revealed the presence of acetone.

Ir analysis of a mixture of equimolar amounts of isopropenyl acetate and methanesulfonic acid in methylene chloride also suggests the formation of mixed anhydride ( $\nu_{\text{max}}$  1800  $\text{cm}^{-1}$  vs. 1825  $\text{cm}^{-1}$  for acetic anhydride). No acetyl methanesulfonate was isolated, however.

This work is part of an extensive study of low temperature acylation procedures using isopropenyl esters in the presence of strong oxy acids. In these applications the mixed anhydride generally is formed *in situ*. Since (1) addition of a stoichiometric quantity of the strong acid is essential for complete acylation to occur, (2) neither the stearic acid-strong acid combination nor the stearic anhydride-strong acid combination is capable of acylation at such ambient temperatures, at reasonable rates, (3) the formation of a mixed anhydride is demonstrated, and (4) the mixed anhydride is shown to be a potent acylating agent, we are confident that the active materials in these ambient temperature acylation reactions are mixed anhydrides. The results of the application studies using the mixed anhydride acylating agents will be published separately.<sup>8</sup>

(1) (a) F. Effenberger and G. Epple, *Angew. Chem., Int. Ed. Engl.*, **11**, 299 (1972). (b) A. Baroni, *Atti Acad. Lincei*, **17**, 1081 (1933). (c) C. G. Overberger and E. Sarlo, *J. Amer. Chem. Soc.*, **85**, 2446 (1963).

(2) G. A. Olah and S. J. Kuhn, *J. Org. Chem.*, **27**, 2667 (1962).

(3) M. H. Karger and Y. Mazur, *J. Amer. Chem. Soc.*, **90**, 3878 (1968).

(4) W. L. Garbrecht, *J. Org. Chem.*, **24**, 368 (1959).

(5) H. J. Hagemeyer, Jr., and D. C. Hull, *Ind. Eng. Chem.*, **41**, 2920 (1949).

(6) E. A. Jeffery and D. P. N. Satchell, *J. Chem. Soc.*, 1876, 1887, 1906 (1962).

(7) E. S. Rothman and S. Serota, *J. Amer. Oil Chem. Soc.*, **48**, 373 (1971).

(8) Detailed experimental data will appear following these pages in the microfilm edition of this volume of the journal. Single copies may be obtained from the Business Operations Office, Books and Journals Division, American Chemical Society, 1155 Sixteenth St., N.W., Washington, D.C. 20036, by referring to code number JOC-73-174. Remit check or money order for \$3.00 for photocopy or \$2.00 for microfiche.

(9) Agricultural Research Service, U.S. Department of Agriculture.

(10) National Research Council-Agricultural Research Service Postdoctoral Research Associate, 1970-1972. Presently Fats and Proteins Research Foundation Fellow.

EASTERN REGIONAL RESEARCH  
LABORATORY<sup>9</sup>  
PHILADELPHIA, PENNSYLVANIA 19118

WILLIAM N. MARMER\*<sup>10</sup>  
SAMUEL SEROTA  
GERHARD MAERKER

RECEIVED SEPTEMBER 22, 1972