

Membrane-Degrading Enzymes in the Tubers of Various Cultivars of *Solanum tuberosum*

Robert A. Moreau

Tubers of 41 potato cultivars were surveyed for phospholipase and *p*-nitrophenyl palmitate hydrolase activities. Phospholipase levels ranged from 2.2 to 29.6 $\mu\text{mol min}^{-1}$ (g of fresh weight) $^{-1}$ and *p*-nitrophenyl palmitate hydrolase ranged from 0.7 to 16.3 $\mu\text{mol min}^{-1}$ (g of fresh weight) $^{-1}$. There was no apparent correlation between the two enzymatic activities among the cultivars tested, thus indicating that the popular *p*-nitrophenyl palmitate hydrolase assay is not an accurate assessment of overall lipolytic activity in potato tubers. A European cultivar, Désirée, which had previously been singled out because of its low levels of lipolytic activity, was found to contain as much phospholipase and *p*-nitrophenyl palmitate hydrolase activities as many of the other cultivars.

Galliard (1970) was the first to report that homogenization of potato tubers resulted in a rapid enzymatic

Eastern Regional Research Center, Agricultural Research Service, U.S. Department of Agriculture, Philadelphia, Pennsylvania 19118.

breakdown of endogenous phospholipids and galactolipids. Even at 0 °C, 26% of the lipids were hydrolyzed after 10 min. At least three different acyl hydrolases have been identified in potato tubers (Galliard, 1971; Hasson and Laties, 1976a; Hirayama et al., 1975; Shepard and Pitt, 1976). One European cultivar, Désirée, was reported to

contain much lower levels of acyl hydrolase than other known cultivars (Galliard and Matthew, 1973). When isolating organelles or membranes from potato tubers, it has therefore been suggested that Désirée should be the cultivar of choice.

In this report 40 common North American cultivars were surveyed for lipolytic enzyme activity and compared with Désirée tubers from two U.S. Department of Agriculture collections. This was done to determine whether any commercially available cultivars had acyl hydrolase activity that was as low or lower than that of Désirée. All samples were tested for their capacity to degrade both a natural membrane lipid (phosphatidylcholine) and a popular synthetic substrate (*p*-nitrophenyl palmitate).

EXPERIMENTAL SECTION

Materials. Potato tubers (*Solanum tuberosum*) of each cultivar were harvested in Maine in Sept 1982 and generously provided by R. Webb, U.S. Department of Agriculture, Agricultural Research Service, Beltsville, MD. In addition, separate samples of Désirée tubers were kindly provided by R. Ross, U.S. Department of Agriculture, Agricultural Research Service, Sturgeon Bay, WI. All tubers were stored at 4 °C for 6 months.

Phosphatidylcholine (dipalmitoyl-¹⁴C) was obtained from New England Nuclear. All other reagents were obtained from Sigma.

Enzyme Preparation. Cores (1-cm diameter) were taken from tubers with a brass cork borer and rinsed in distilled water. Samples (4 g) were cut from the cortex and homogenized in a semimicro Waring Blendor in 40 mL of 0.1 M potassium phosphate buffer (pH 7.0) and 2 mM sodium metabisulfite at 0 °C. The homogenate was filtered through two layers of cheesecloth and centrifuged at 20000g for 30 min. The supernatant was collected and enzyme activities were measured within 4 h. Both enzyme activities were stable at 0 °C for at least 4 h.

Phospholipase Assay. Phospholipase activity was routinely measured by using [¹⁴C]phosphatidylcholine (PC). The substrate was prepared by sonicating labeled PC (dipalmitoyl-¹⁴C) and unlabeled PC in 0.4% Triton X-100 in a bath sonicator (Bransonic 12, Branson Inc., Shelton, CT) until the emulsion became crystal clear. The standard reaction mixture (1 mL) contained 1 mM PC-dipalmitoyl (50 000 dpm), 0.2% Triton X-100, 100 mM Tricine-NaOH (pH 8.5), 1 mM CaCl₂, and 2–5 μL of enzyme. Two different enzyme concentrations were tested for each cultivar. The reaction vials were incubated in a shaking water bath (140 rpm) at 30 °C for 30 min. The enzymatic reactions were simultaneously stopped and extracted by adding 50 μL of acetic acid, 7 mL of 3:2 hexane–2-propanol (v/v), and 5 mL of 6.7% Na₂SO₄. The lipids were removed in the top phase and dried under a stream of N₂. Fifty micrograms each of oleic acid and PC (soybean) was added as carriers, and the lipid samples were spotted on 250-μm silica gel G TLC plates and developed in 70:30:1.5 hexane–diethyl ether–acetic acid (v/v/v). The free fatty acid and PC spots were visualized with I₂. The radioactivity in the two regions was measured with a Berthold LB 283 linear analyzer. The percent conversion of [¹⁴C]PC to ¹⁴C free fatty acid was used to calculate enzyme activity based on the initial amount of substrate (1 μmol).

Esterase Activity. The activity of *p*-nitrophenyl palmitate hydrolase (PNP hydrolase) was measured spectrophotometrically as described by Galliard and Matthew (1973). The reaction mixture (1 mL) contained 0.40 mM PNP, 0.2% Triton X-100, 50 mM potassium phosphate (pH 8.0), and 2–20 μL of enzyme. The change

in absorbance at 405 nm was measured continuously for 3–4 min with a Beckman Model 35 spectrophotometer. Three different enzyme concentrations were tested for each cultivar.

Protein Assay. Samples of tuber supernatants were mixed with an equal volume of 10% trichloroacetic acid and incubated at 0 °C for 18 h. The mixture was then centrifuged at 5000g for 10 min. The pellet was resuspended in 0.1 N NaOH, and protein was assayed by the Lowry–Markwell procedure (Markwell et al., 1978).

RESULTS

Preliminary experiments were conducted in order to determine the optimal conditions for assaying phospholipase activity. By use of a homogenate from Katahdin tubers, maximum phospholipase activity was obtained with 1 mM PC, 0.2% Triton X-100, 100 mM Tricine-NaOH (pH 8.5), and 1 mM CaCl₂. Because these conditions were very similar to those of two previous studies (Hasson and Laties, 1976b; Hirayama et al., 1975), phospholipase activity in all cultivars was measured under these conditions.

Among the 41 cultivars that were obtained from Maine, total phospholipase activity ranged from 2.20 to 29.60 μmol min⁻¹ (g of fresh weight)⁻¹ and the specific activity ranged from 0.94 to 6.43 μmol min⁻¹ (mg of protein)⁻¹ (Table I). PNP hydrolase activity ranged from 0.66 to 16.32 μmol min⁻¹ (g of fresh weight)⁻¹. There was no apparent correlation between PNP hydrolase and phospholipase activities. Because the PNP hydrolase and phospholipase activity in Désirée tubers was much higher than previously reported (Galliard and Matthew, 1973), a second batch of Désirée tubers was obtained from Sturgeon Bay, WI. The Désirée tubers from the Wisconsin collection had phospholipase and PNP hydrolase activities that were very similar to those in the tubers from the Maine collection.

The concentration of soluble protein ranged from 2.05 to 4.92 mg/g of fresh weight. In general, the cultivars with the lowest total phospholipase activity also had the lowest soluble protein levels, and the higher phospholipase cultivars had higher soluble protein levels.

DISCUSSION

The purpose of this study was to compare the levels of membrane-degrading enzymes in the tubers of commercially available cultivars with those of Désirée. These results indicate that, at least for the North American scientist, there is no advantage in choosing Désirée tubers over any of the North American cultivars. Among the cultivars that were surveyed, Russett Burbank had the lowest phospholipase levels. Because it has the lowest phospholipase content and because it is the most popular North American cultivar, Russett Burbank should probably be the cultivar of choice when membrane destruction needs to be minimized. However, even a tuber homogenate of Russett Burbank has the potential ability (at pH 8.5 and 30 °C) to degrade all of its membrane lipids in less than 1 min [based on an average lipid content of 1 mg/g of fresh weight as measured by Galliard (1973)]. It is therefore obvious that cultivar selection is only the first of several precautions that need to be followed to minimize membrane breakdown. Working at 0 vs. 25 °C is helpful, but it only reduces membrane hydrolysis by about 60% (Galliard, 1970; Hasson and Laties, 1976b). Freshly harvested tubers (Wardale, 1980) or tubers stored at 3 vs. 7 °C (Walcott et al., 1982) may be preferable because they contain less PNP hydrolase activity. Some organelles such as mitochondria can be sedimented fast enough to minimize membrane destruction (Hasson and Laties, 1976b). However, other subcellular fractions such as plasma membranes, endoplasmic reticulum, and Golgi vesicles

Table I. Levels of Phospholipase and *p*-Nitrophenyl Palmitate Hydrolase in Potato Tubers^a

| cultivar | phospholipase | | PNP hydrolase, $\mu\text{mol min}^{-1}$ (g of fresh wt) ⁻¹ | protein, mg (g of fresh wt) ⁻¹ |
|---------------------|---|--|---|--|
| | $\mu\text{mol min}^{-1}$ (g of fresh wt) ⁻¹ | $\mu\text{mol min}^{-1}$ mg ⁻¹ | | |
| Russett Burbank | 2.20 | 0.94 | 2.31 | 2.35 |
| Hudson | 2.63 | 1.18 | 0.80 | 2.23 |
| Green Mountain | 3.84 | 1.87 | 0.66 | 2.05 |
| Red La Sada | 3.86 | 1.17 | 5.57 | 3.30 |
| Lemki | 4.01 | 1.51 | 2.14 | 2.66 |
| Cobbler | 4.27 | 1.37 | 3.59 | 3.12 |
| Désirée (Wisconsin) | 4.62 | 1.26 | 2.42 | 3.67 |
| Désirée (Maine) | 5.69 | 1.93 | 1.77 | 2.95 |
| Bounty | 5.87 | 2.41 | 1.59 | 2.44 |
| Red Pontiac | 6.27 | 1.97 | 3.62 | 3.19 |
| Centennial | 6.44 | 2.45 | 2.50 | 2.63 |
| Butte | 6.85 | 1.77 | 3.73 | 3.87 |
| White Rose | 7.64 | 3.07 | 1.74 | 2.49 |
| Houma | 7.95 | 2.25 | 3.14 | 3.54 |
| Abnaki | 8.01 | 2.86 | 3.06 | 2.80 |
| Crystal | 8.52 | 2.38 | 4.29 | 3.58 |
| Bison | 9.31 | 2.76 | 2.70 | 3.37 |
| Campbell-12 | 9.44 | 2.83 | 8.34 | 3.34 |
| Saco | 10.09 | 3.01 | 7.97 | 3.35 |
| Campbell-13 | 10.15 | 2.48 | 3.20 | 4.09 |
| La Rouge | 10.21 | 2.27 | 12.21 | 4.49 |
| Buckskin | 10.45 | 3.49 | 4.19 | 2.99 |
| Katahdin | 10.88 | 3.60 | 1.75 | 3.02 |
| Superior | 12.64 | 3.39 | 10.70 | 3.73 |
| Dakchip | 12.86 | 3.38 | 10.36 | 3.81 |
| Kennebec | 13.06 | 3.80 | 7.23 | 3.44 |
| Wausseon | 13.27 | 3.51 | 2.68 | 3.78 |
| Bel Rus | 13.36 | 3.51 | 6.74 | 3.81 |
| Sebago | 13.48 | 4.29 | 8.16 | 3.14 |
| Norgold Russett | 13.70 | 4.44 | 16.32 | 3.08 |
| Alaska Red | 13.74 | 3.56 | 2.75 | 3.86 |
| Campbell-11 | 15.16 | 3.85 | 3.93 | 3.94 |
| Norchip | 15.32 | 4.17 | 5.85 | 3.67 |
| Atlantic | 16.93 | 3.59 | 4.05 | 4.72 |
| Monoma | 17.08 | 4.80 | 6.79 | 3.56 |
| Ontario | 17.36 | 4.47 | 6.49 | 3.88 |
| Denali | 20.25 | 4.98 | 4.03 | 4.07 |
| Belchip | 20.47 | 4.26 | 3.93 | 4.80 |
| Pungo | 22.62 | 5.76 | 11.38 | 3.93 |
| La Chipper | 23.91 | 5.45 | 6.09 | 4.39 |
| Cherokee | 25.85 | 5.25 | 8.02 | 4.92 |
| Highlat | 29.60 | 6.43 | 9.34 | 4.60 |

^aData represent the averages of two or three separate experiments (separate tuber samples, homogenizations, and enzyme assays) for each cultivar. The deviations from the mean were less than 30% for all cultivars.

require longer centrifugation times and are therefore nearly impossible to isolate from potato tubers without extensive membrane degradation. Another alternative is to add exogenous inhibitors to the homogenizing medium. Inhibitors such as bovine serum albumin (BSA) (Hasson and Laties, 1976b), diisopropyl fluorophosphate (Galliard, 1971; Matsuda and Hirayama, 1979), and dibucaine (Pun et al., 1980) have been reported to be effective.

PNP hydrolase activity in the Désirée tubers of this study was about 20 times higher than that reported by Galliard and Matthew (1973) or Wardale (1980). The reason for this discrepancy is not clear. Perhaps differences in cultural practices or nutrient availability in Great Britain and in the United States of America may account for this observation. Because this clone was introduced into the United States directly from Great Britain, the genotype should be unchanged. The levels of PNP hydrolase and phospholipase in the 40 other cultivars are consistent with those previously reported (Galliard and Matthew, 1973; Hasson and Laties, 1976b; Matsuda and Hirayama, 1979). The level of phospholipase activity in Désirée tubers has not previously been published; however, a recent report (Theologis and Laties, 1980) demonstrated similar rates of phospholipid breakdown in tuber slices of Désirée and Russett Burbank (12% vs. 19% loss of lipid

phosphorus per 2 h, respectively).

There was no apparent correlation between phospholipase activity and PNP hydrolase activity in the 41 cultivars. This is consistent with the results of Shepard and Pitt (1976), who were able to separate phospholipase activity and PNP hydrolase activity by isoelectric focusing. These results indicate that screening potato tubers with the convenient PNP hydrolase assay used in the previous survey (Galliard and Matthew, 1973) is not an accurate assessment of overall lipolytic activity.

ACKNOWLEDGMENT

I thank A. Baker for excellent technical assistance and Dr. R. Webb and R. Ross for supplying tubers.

Registry No. PNP-hydrolase, 9001-62-1; phospholipase, 9013-93-8.

LITERATURE CITED

- Galliard, T. *Phytochemistry* 1970, 9, 1725.
 Galliard, T. *Biochem. J.* 1971, 121, 379.
 Galliard, T. *J. Sci. Food Agric.* 1973, 24, 617.
 Galliard, T.; Matthew, J. A. *J. Sci. Food Agric.* 1973, 24, 623.
 Hasson, E. P.; Laties, G. G. *Plant Physiol.* 1976a, 57, 142.
 Hasson, E. P.; Laties, G. G. *Plant Physiol.* 1976b, 57, 148.
 Hirayama, O.; Matsuda, H.; Takeda, H.; Maenaka, K.; Takatsuka, H. *Biochim. Biophys. Acta* 1975, 384, 127.

Markwell, M. A. K.; Haas, S. M.; Bieber, L. L.; Tolbert, N. E. *Anal. Biochem.* 1978, 87, 206.
Matsuda, H.; Hirayama, O. *Bull. Fac. Agric., Shimane Univ.* 1979, 13, 105.
Pun, W. H.; Khan, A. A.; Chung, I.; Haydar, M.; Hadziyev, D. *Potato Res.* 1980, 23, 57.
Shepard, D. V.; Pitt, D. *Phytochemistry* 1976, 15, 1471.
Theologis, A.; Laties, G. G. *Plant Physiol.* 1980, 66, 890-896.
Walcott, P. J.; Kenrick, J. R.; Bishop, D. G. In "Biochemistry and

Metabolism of Plant Lipids"; Wintermans, J. F. G. M.; Kuiper, P. J. C., Eds.; Elsevier: Amsterdam, 1982; p 297.
Wardale, D. A. *Phytochemistry* 1980, 19, 173.

Received for review June 8, 1984. Accepted October 2, 1984.
Reference to brand or firm name does not constitute endorsement
by the U.S. Department of Agriculture over others of a similar
nature not mentioned.